

Simulation of rice production under climate change scenarios in the Southern coasts of Caspian Sea

H. Mohammadi^{a*}, Gh. Azizi^a, D. Mazaheri^b, F. Rabbani^a

^a Natural Geography Department, Faculty of Geography, University of Tehran, Tehran, Iran

^b Faculty of Agricultural Science and Engineering, University of Tehran, Karaj, Iran

Received: 25 April 2015; Received in revised form: 2 June 2015; Accepted: 14 June 2015

Abstract

Climate change has direct and indirect consequences on crop production and food security. Agriculture and crop production is one of the factors which depend on the weather conditions and it provides the human requirements in many aspects. The objective of this study is to assess the impacts of future climatic change on irrigated rice yield using the CERES-Rice model in the Southern Coast of Caspian Sea under three climate change scenarios of Sra1b, Sra2 and Srb1. Required data for this research includes the meteorological, soil and crop management data. The meteorological data include the daily data of minimum temperature, maximum temperature, solar radiation and precipitation during 1981-2010 and Global Climate Models (HADCM3, ECHAM5, IPCM4, GFCM2, NCCCSM and INCM3) during 1971-2000. Soil and product management data provided from field experiment was conducted from 2008 to 2009 at the Rice Research Institute in Rasht. Validating of Global Climate Models show that ECHAM5 climate model has the highest correlation with the lowest error to simulate the future temperature and precipitation. We used ECHAM5 climate model coupled with a crop growth model for simulating of the effects of climate change on rice production. The results of prediction of climate change scenarios show that minimum and maximum temperature will be ascending and precipitation will be decreasing in the Rasht station. Results of simulated yield and biomass of the rice crop base on scenarios of Sra1b, Sra2 and Srb1 show that rice crop yield and biomass decrease with increasing of mean temperature and decreasing of precipitation.

Keywords: Climate change; CERES-Rice model; ECHAM5; Southern Coast of Caspian Sea; Rasht

1. Introduction

Climate Change is the most important threat for water resources, soil, agriculture, human health and food security. Food security is one of the main goals of the government's agenda. The role of agriculture due to its contribution to promote the economic growth and development and food security for a growing population, production, exchange, employment and development activities related to crop production is important from the perspective of economic development. On the other hand, agriculture and crop production is one of the factors which depend on the weather conditions and it provides the human requirements in

many aspects. Future climate change can have considerable effects on growth, performance and water consumption for crops. If it happens very fast, then the plants will take the most damage (Bazzaz and Sombroek, 1996). In general, organisms such as plants have adapted to their environment over a long period of time, and had matched their growth with this condition. In this regard, any rapid change in weather conditions will cause significant changes in the patterns of growth and development of plants and depending on the changes; the plants will have spatial and temporal displacement and ultimately may even exclude them from the agricultural system in a particular area (Horie *et al.*, 2000).

The increasing atmospheric CO₂ and other greenhouse gases due to anthropogenic are well documented and theoretical reasons for higher concentrations of these gases to cause global

* Corresponding author. Tel.: +98 912 3849767,
Fax: +98 61113536.
E-mail address: hmmohammadi@yahoo.com

warming in recent years. The fourth assessment report (AR4) of the IPCC indicates that the atmospheric CO₂ is accumulating in the atmosphere faster than past and global temperatures have increased in the past 100 years by an average of 0.74°C (IPCC, 2007). Evidences show the maximum and minimum temperatures are likely to increase at different rates, often with a faster increase in minimum temperature than maximum temperature (Easterling *et al.*, 1997). Rainfall in the high latitudes of the northern hemisphere has increased, while rainfall in eastern Asia, Australia, the Sahel and the Pacific region has declined, with rainfall variability increasing almost everywhere in the world (Dore, 2005). The changing of atmospheric concentration of CO₂, increasing temperatures and reduced solar radiation and increased ozone concentrations, increasing of rainfall variability, severe and frequent droughts have serious direct and indirect consequences on crop production and food security (Reddy and Hodges, 2000).

Many researches have been conducted to determine the effects of climate change and other environmental factors on growth and yield of crops over the past two decades. Auffhammer *et al.* (2012) studied impact the monsoon rainfall change in India using Monte Carlo simulation. They concluded that yield would have received an additional boost of nearly 4% if warmer nights and lower rainfall at the end of the growing season had not occurred. Mainuddin *et al.*, (2013) studied impact of climate change on rice in the lower Mekong Basin base on ECHAM4 global climate model and A2 and B2 scenarios. The results show that yield of rain fed rice may increase in the upper part of the basin in Laos and Thailand and may decrease in the lower part of the basin in Cambodia and Vietnam. Gohari *et al.*, (2013) evaluated climate change impacts on crop production and water productivity of four major crops (wheat, barley, rice, and corn) in Iran's Zayandeh-Rud River Basin. They show crop production and water productivity of all crops is expected to decrease due to lower precipitation and higher water requirements under higher temperature. Soora *et al.* (2013) assessed the regional vulnerability of rice to climate change in India using A1b, A2, B1 and B2 emission scenarios and global climate model (MIROC3.2.HI). The study projected a progressive reduction in irrigated rice yields due to climate change towards the end of the century, if no adaptation is followed; while for rain fed rice, negative impacts likely to reduce with time due to projected increase in

rainfall in many areas. Poudel and Kotani (2013) studied climatic impacts on crop yield and its variability in Nepal. The results show that an increase in the variance of both temperature and rainfall has adverse effects on crop productions in general. On the other hand, a change in the mean levels of the temperature and rainfall induces heterogeneous impacts, which can be considered beneficial, harmful or negligible, depending on the altitudes and the kinds of crops.

The results, depending on the type of plant photosynthetic pathway and geographic area, are varied and having an overall statement about the response of different species to climate change needs to conduct a case study.

Rice is a main food for half of the world's population. In Iran, rice is the second important staple food crop after wheat and the main rice cultivation center is across of plains in southern coasts of Caspian Sea. This region provides 70% of total rice production areas in Iran. (Iran Ministry of Agriculture, 1993).

The objective of this study is to assess the impacts of future climatic change on irrigated rice yield in the Caspian coastal area under three climate change scenarios of SRA1B, SRA2 and SRB1. Different instruments must be used to predict or simulate the effects of climate change on crops so that the best policy making and planning can be conduct and ultimately provide the food security and the maximum economic welfare of manufacturers.

2. Materials and Methods

2.1. Study area

The Rasht station is selected as the rice cultivation center in the southern coasts of Caspian Sea with long-time weather data, to simulate the effects of climate change on rice yield in this study. Mean annual precipitation is 111.5 mm and maximum rainfall occurs in autumn. The mean maximum air temperatures ranged between 11 to 30 °C and the mean minimum air temperatures is varied between 6 to 26 °C. Summer with humid and warm climate and low rainfall is suitable for the long growth of rice.

2.2. Model input data

Required data for this research includes the meteorological, soil and crop management data. Required meteorological data include the collection of daily data of Rasht stations (minimum temperature, maximum temperature,

solar radiation and precipitation) during 1981-2010 and Global Climate Models (HADCM3, ECHAM5, IPCM4, GFCM2, NCCCSM and INCM3) during 1971-2000 period. Soil and crop management data have provided from field experiments. Soil data include the classification of the soil, the water-holding characteristics of different soil layers plus their bulk density, organic C, pH, drainage coefficient and root growth factor. Initial values of soil water, nitrate, and ammonium are needed as well as an estimate of the above and belowground residues from the previous crop.

Crop-management factors include planting date, planting depth, row spacing and direction, plant population, fertilization, irrigation, inoculation, residue applications, tillage, and harvest date.

2.3. Generating of Future Weather data

The future climate change scenarios that are usually generated using global circulation models cannot provide the details on very small spatial scales due to scientific limitations and incomplete observational data (IPCC, 2007). For decreasing of gap between the scale of GCMs and required resolution for practical applications, downscaling provides climate change information at a suitable local scale from the GCM data (Hewitson and Crane, 1996).

We used Lars- Weather Generator model for simulating time-series of daily weather and downscaling climate models at a single site in this study. It was calibrated and validated using statistical tests between generated and observed data. Although the Lars-Wg model contains 15 global climate models, the six models (HADCM3, ECHAM5, IPCM4, GFCM2, NCCCSM and INCM3) have three scenarios SRA1B, SRA2 and SRB1. In evaluating the accuracy of the atmosphere general circulation models in production of future weather data, we validated six global climate models using the absolute root mean square error (RMSE) and mean absolute error (MAE) between the simulated and observed data. Future weather data generated for 2011-2032 and 2046- 2065 base on climate change scenarios.

2.4. Field Experiment

Field experiment was conducted from 2008 to 2009 at the Iranian Rice Research Institute in Rasht (37 12 N, 49 38 E) during the rice cultivation season. The rice crop varieties Hashemi and Alikazemi have been selected for

mode calibration that widely cultivated in the north of Iran during the spring season.

The experiment was done in a split-plot design with three irrigation regimes as the main plot, four N levels and three replications. The plot size for the subplots was 15 m². The irrigation regimes were continuous submergence with irrigation at 5-day intervals and 8-day interval. The four N rates applied were 0, 30, 60, and 90 kg N ha⁻¹. The subplots (15 m²) consisted of four N levels: N1: no N application; N2: total N rate of 45 kg ha⁻¹; N3: total N rate of 60 kg ha⁻¹; and N4: total N rate of 75 kg ha⁻¹.

Seedlings were grown in wet beds for approximately 25–30 days and was done transplanting at three plants per hill with a spacing of 20 × 20 cm. seeding was done in the nursery in early April. Thirty-day-old seedlings were transplanted in early May. Treatments were harvested in mid August. Weeds, insects, and diseases were controlled in all plots to avoid yield loss.

Soil physical and chemical properties such as texture, bulk density, hydraulic conductivity, drained upper limit, drained lower limit, field capacity, pH, cation-exchange capacity (or CEC that is the total capacity of a soil to hold exchangeable cations), organic carbon (OC), total N, phosphorus (P), potassium (K) were determined up to a depth of 80 cm, at an interval of 10 cm, following standard procedures (Table 1).

2.5. Crop Model Description

The crop model that used for simulating rice response to climate change is Ceres-rice model. The Crop Estimation Resource and Environment Synthesis (CERES) - Rice model was one of the DSSAT models developed by the International Benchmark Sites Network for Agrotechnology Transfer (IBSNAT) project which can simulate growth, development and yield of rice varieties by numerical integration of constituent processes under different agroclimatic conditions and management strategies (Mathauda *et al.*, 2000, Matthews *et al.*, 2002). Ceres-Rice has been evaluated for many tropical and sub-tropical locations in Asia and in temperate climates (Timsina and Humphreys, 2006).

Databases contain weather, soil, experiment conditions and measurements, and genotype information for applying the models to different situations. The main function of the weather module is to read or generate daily weather data. It reads in daily weather values

(maximum and minimum air temperatures, solar radiation and precipitation, relative humidity, and wind speed when available) from the daily weather file. The soil in the land unit is represented as a one-dimensional profile; it is homogenous horizontally and consists of a number of vertical soil layers. The soil module integrates information from four submodules: soil water, soil temperature, soil C and N, and soil dynamics. The soil-plant-atmosphere module computes daily soil

evaporation and plant transpiration. According to Jones et al (2003) this software helps users prepare these databases and compare simulated results with observations to give them confidence in the models if modifications are needed to improve accuracy. In addition, programs contained in DSSAT allow users to simulate options for crop management over a number of years to assess the risks associated with each option (Jones *et al.*, 2003).

Table 1. Physical and chemical per soil layers of the experiment field

| Soil properties | Depth(CM) | | | | | |
|-----------------------------|-----------|-------|-------|-------|-------|-------|
| | 0-10 | 10-20 | 20-30 | 30-40 | 40-60 | 60-80 |
| Clay (%) | 47 | 44 | 47 | 47 | 49 | 61 |
| Sand (%) | 14 | 17 | 9 | 11 | 9 | 5 |
| Loam (%) | 39 | 39 | 44 | 42 | 42 | 34 |
| Bulk density (g cm) | 1.10 | 1.20 | 1.32 | 1.31 | 1.33 | 1.29 |
| Water content at saturation | 0.65 | 0.62 | 0.62 | 0.60 | 0.60 | 0.60 |
| Water content at FC | 0.40 | 0.40 | 0.41 | 0.42 | 0.42 | 0.39 |
| Water content at PWP | 0.27 | 0.30 | 0.30 | 0.30 | 0.32 | 0.29 |
| Ksat | 57.5 | 30.8 | 0.4 | 11.4 | 10.4 | 21.4 |
| PH | 7.1 | 7.23 | 7.26 | 7.08 | 7.5 | 7.5 |
| CEC (meq/100g-1) | 33 | 32 | 31 | 31 | 30 | 30 |
| Organic carbon (%) | 0.16 | 0.14 | 0.074 | 0.074 | 0.09 | 0.09 |
| P (ppm) | 10.1 | 7.3 | 5.2 | 3.2 | 2.1 | 1.7 |

Phosphorus (P) and potassium (K) were applied at transplanting in all plots at the rate of 25 kg phosphorus (P) ha⁻¹ and 75 kg potassium (K) ha⁻¹. The dates of emergence, panicle initiation, flowering, and physiological maturity were recorded for each management.

2.6. Crop Model Calibration and Validation

The Ceres-Rice model has been evaluated separately for the calibration data set of 2008 and for the validation data set of 2009. Calibration is defined as a set of operations that adjust some model parameters, under specified conditions, to local conditions and it creates genetic coefficients for new cultivars. The calibration of the Ceres-Rice model was based on data from end-of-season samplings of grain yield, crop biomass, and N contents in biomass and in grain in 2008 field experiment. Growth and development of crop varieties in maturity are distinguished using the genetic coefficients in Ceres-rice models. The

genetic coefficients of the cultivar of Hashemi and Alikazemi that affect the occurrence of phenological stages in the Ceres-Rice models were derived using the Genotype Coefficient Calculator program (GENCALC) which is part of the Decision Support System for Agrotechnology Transfer (DSSAT). This program estimates the coefficients for a genotype by iteratively running the crop model with an approximate value of the coefficients concerned. It alters the cultivar coefficients automatically until the simulated and measured values match (Hunt et al., 1993). Table 2 shows results of calibrating the Ceres - Rice model in 2008 field experiment for cultivars of Hashemi and Alikazemi.

Table 2. Genetic coefficients of the rice cv. Hashemi and Alikazemi as derived by GENCALC of DSSAT model

| Genetic parameters | Hashemi | Alikazemi |
|--------------------|---------|-----------|
| P1 | 310 | 320 |
| P2R | 20 | 20 |
| P5 | 350 | 360 |
| P20 | 13.5 | 13.5 |
| G1 | 55 | 55 |
| G2 | 0.025 | 0.028 |
| G3 | 1 | 1 |
| G4 | 1 | 1 |

P1 is respectively thermal time in the basic vegetative phase of the plant, P20 is critical photoperiod of development occurs at a maximum rate, P2R is photoperiod sensitivity in panicle initiation, P5 is grain filling duration, G1 is potential spikelet number coefficient, G2 Single grain weight under ideal growing conditions, G3 tillering coefficient under ideal conditions, G4 is temperature tolerance coefficient.

We compared the simulated and measured data for validating of model to increase confidence in the ability of model. The combination of graphical and statistical methods was used to compare the simulated and observed results (final biomass and yield). So, we evaluated model performances using the absolute root mean square error (RMSE), root mean square error normalized (RMSEn) and mean absolute error (MAE). RMSE and RMSEn are well statistical methods to test the goodness of fit of simulation models (Bouman and Van Laar, 2006). Willmott and Matsuura (2005) suggested that MAE is the most natural measure of average error magnitude and is an unambiguous measure of average error magnitude. Willmott *et al.*, (1985) suggested some statistical methods for evaluating of model and offered that RMSE is the “best” measure as it summarizes the mean difference in the units of observed and simulated values:

$$RMSEa = [1/n \sum (Yi - Xi)^2]^{0.5} \tag{1}$$

$$RMSEn = 100. [RMSEa / X_{obs}] \tag{2}$$

$$MAE = \sum |Yi - Xi| / n \tag{3}$$

Where Yi is the simulated value, Xi is the measured value, and n is the number of measurements. It was calculated the slope of linear relation between simulated and observed values (), intercept (), and coefficient of determination (R²) of the linear regression between simulated (Xsim) and measured (Xobs) values For the same variables. We also calculated the Student’s t-test of means assuming unequal variance (P (t*)) between simulated and measured values. Scatter plots between simulated and measured data were used to show the overall goodness-of-model fit.

2.7. Simulation of Rice Yield and Biomass

Crop simulation models are an efficient tool for studying the present and future effects of climate changes on crop. Because of the limitations and weak of statistical models, the simulation of the effects of climate change on agriculture usually employs a climate model coupled with a crop growth model (Yao et al., 2011). We simulated

the yield and biomass of rice base on the prediction of GCM under climate change scenario Sra1b, Sra2 and Srb1 during 2011-2032 and 2046-2065. The stages of performance of model are including as follows:

- defining the weather data (minimum and maximum temperature (°C), Solar Radiation (MJ/m2) and rainfall (mm)). For utilizing of weather data in DSSAT, it should be formatted to fit the system. Weatherman is the program for introducing climate data for using in crop simulation in DSSAT.
- creating a new experiment and input the set of crop management data base on field experiment (planting and harvesting date, choosing cultivars, defining amount and timing of fertilization and irrigation, soil, weather, treatments and simulation option).
- computing of changes in crop productivity as result of climate change.

3. Results and Discussion

3.1. Downscaling and Climate Model Evaluation

The Lars-WG model was validated using statistical tests such as the Kolmogorov –Smirnov (K–S) test, T- test and F-test with P-value. The K–S test was done for testing equality of the seasonal distributions of wet and dry series (WDSeries), distributions of daily rainfall (RainD), and distributions of daily minimum (TminD) and maximum temperature (TmaxD). The T-test is performed on testing equality of monthly mean rainfall (RMM), monthly mean of daily maximum temperature (TmaxM), and monthly mean of daily minimum temperature (TminM) and The F-test is performed on testing equality of monthly variances of precipitation (RMV) calculated from observed data and downscaled data. If the P- value of a data set is below a certain pre-determined amount (0.05), the "null hypothesis" of their experiment rejects. The null hypothesis is that the simulated data is same observed data. Table 3 shows the p-value of statistical test for validating of the Lars-WG model in Rasht stations.

Table 3. The P-value of statistical test for validating of the Lars-WG model

| Statistical Test | WDSeries | RainD | RMM | RMV | TminD | TminM | TmaxD | TmaxM |
|------------------|----------|-------|--------|--------|-------|--------|-------|--------|
| | K-S | K-S | T test | F test | K-S | T test | K-S | T test |
| winter | 0.79 | 1 | 0.5 | 0.35 | 0.99 | 0.55 | 0.99 | 0.34 |
| spring | 1 | 0.89 | 0.59 | 0.34 | 1 | 0.33 | 1 | 0.63 |
| autumn | 1 | 1 | 0.79 | 0.74 | 0.99 | 0.45 | 0.99 | 0.55 |
| summer | 1 | 0.99 | 0.67 | 0.2 | 0.99 | 0.62 | 1 | 0.74 |

The global climate models validated using measures of accuracy indicators during baseline period (1981-2000). The measures of accuracy indicators show that ECHAM5 climate model (European Community Hamburg atmospheric model coupled to a large scale geostrophic ocean model that was generated by Max-Planck Institute for Meteorology in Germany) has the highest

correlation with the lowest error to simulate the temperature and precipitation (Table 4). Therefore, according to the uncertainty principle, due to the low performance of other models in simulating the temperature and precipitation for reducing the errors in simulating, they won't be used in simulation of future climate.

Table 4. Validation of global climate models using measures of accuracy indicators

| Global Climate Models | RMSE | | MAE | |
|-----------------------|------------------|--------------------|------------------|--------------------|
| | Temperature (°C) | Precipitation (mm) | Temperature (°C) | Precipitation (mm) |
| GFCM2 | 8/5 | 3/34 | 8/35 | 2/81 |
| HADCM3 | 6/06 | 3/61 | 5/34 | 2/99 |
| INCM3 | 5/98 | 3/5 | 5/73 | 2/96 |
| IPCM4 | 4/4 | 3/44 | 3/9 | 2/91 |
| ECHAM5 | 2/69 | 3/19 | 2/36 | 2/69 |
| NCCCSM | 4/28 | 2/96 | 4/09 | 2/53 |

3.2. Crop Model Evaluation

The results of statistical methods used to evaluate the model performance are shown in Tables 5. The RMSE was 306.3 kg ha⁻¹, the normalized RMSE was 1.34 % and the MAE was 270 for rice crop yield. The biomass was slightly over predicted with an RMSE of 480.7 kg ha⁻¹, the normalized RMSE was 0.89% and the MAE was 354.2 for total biomass. Paired t-test showed no significant differences between the measured

and simulated yield and total biomass values at P=0.05 confidence level.

Figure 1 shows the comparison of simulated with measured yield and final biomass for all data of the validation sets. The linear regression between simulated and measured values had a slope close to 1, an intercept that was small (compared with the range in the recorded values), and r relatively close to 1, indicating a close correlation between the simulations and the measurements.

Table 5. Results for CERES rice simulations of final aboveground biomass and yield for the calibration conditions

| year | Crop Parameters | X obs | X sim | | R ² | P(t) | RMSE | RMSEn | MAE | |
|------|--------------------------------|-------|--------|------|----------------|------|------|-------|------|-------|
| 2009 | Biomass (kg ha ⁻¹) | 6704 | 6747.7 | 0.82 | 1238 | 0.94 | 0.5 | 480.7 | 0.89 | 354.2 |
| | Yield (kg ha ⁻¹) | 2843 | 3114 | 0.89 | 570.7 | 0.87 | 0.2 | 306.3 | 1.34 | 270 |

Xobs, Mean of measured values; Xsim, Mean of simulated values; P(t*), Significance of paired T-test at 95%; β , Slope of linear relation between simulated and measured values; α , Intercept of linear relation between simulated and measured values; R², coefficient of determination; RMSEa, Absolute root mean square error; RMSEn, Normalized root mean square error (%); MAE, mean absolute error.

3.3. Estimating Rice Yield and Biomass Under Climate Change Scenarios

It can be concluded that the Lars-WG model has good performance in the Rasht stations in generating daily minimum and maximum air temperature and daily precipitation. So, it was used to predict daily Tmax and Tmin and daily precipitation for the Rasht station during 2011–2032 and 2046–2065 under three scenarios of Sra1b, Sra2 and Srb1. The prediction results of climate variables in the Rasht station show that minimum and maximum temperature will increase in future. Also, the minimum air temperature will increase more than maximum air temperature. The prediction results under three

scenarios of Sra1b, Sra2 and Srb1 in comparison with present level showed that the maximum temperature increase will be 0.7 – 2.0°C during 2011-2032 and 1.5–2.2°C during 2046-2065. The minimum temperature increase will be 0.7–2.2°C during 2011-2032 and 1.6–2.3°C during 2046-2065. The ECHAM5 climate model has predicted the mean precipitation level lower than the mean observation course data on annual precipitation. Annual rainfall will decrease in Rasht station comparison to the normal level and it is related to decreasing of summer rainfall. Also, study of seasonal rainfall showed that rainfall will decrease 7% in the rice growing season (April-august) in comparison to the normal level in the Rasht station (Table 6).




Fig. 1. Evaluating of simulated versus observed yield and Biomass. , Slope of linear relation between simulated and measured values that is close to 1; , Intercept of linear relation between simulated and measured values that is small; r, Correlation Coefficient is close to 1

Table 6. Results of model's predictions under three scenarios of Sra1b, Sra2 and Srb1 in the Rasht station

| Climate variables | Observed period (1981-2010) | 2011-2032 | | | 2046-2065 | | |
|--------------------------|-----------------------------|-----------|-------|-------|-----------|-------|-------|
| | | Sra1b | Srb1 | Sra2 | Sra1b | Srb1 | Sra2 |
| Minimum Temperature (°C) | 11.97 | 14.23 | 12.88 | 12.75 | 14.29 | 13.61 | 14.04 |
| Maximum Temperature (°C) | 20.45 | 22.6 | 21.34 | 21.18 | 22.67 | 22.07 | 22.51 |
| Precipitation (mm) | 3.69 | 3.34 | 3.31 | 3.34 | 3.39 | 3.54 | 3.51 |

The results from ECHAM5 show mean annual temperature increase will be 0.6 –2.2 °C. Higher temperature shortens the crop growth period; consequently reduce the available to the plant for photosynthetic accumulation (Ritchie, 1993). High temperature conditions after heading result in smaller kernels at maturity leading to reduction in grain yield (Yoshida and Hara, 1977). In this study the response of rice crop to changes in atmospheric temperature is studied and presented in Figure 4. The figure shows a decrease in Hashemi and Alikazemi rice yield and biomass

for a 0.6 to 2°C rise in temperature from present day level. At 2°C temperature rise the yield decreases 24% and 27% respectively in Hashemi and Alikazemi rice from the base temperature level. The biomass decreases 26% in Hashemi and Alikazemi from the base temperature level.

Figure 5 shows the rice crop yield and biomass in response to changes in precipitation for 0 to 15 mm drop from present day level. The results show that the decrease in yield and biomass due to decrease in precipitation less than observed values.




Fig. 4. Sensitivity of rice yield and biomass to temperature changes between 0 to 2°C as simulated by the Ceres-Rice model




Fig. 5. Sensitivity of rice yield and biomass to precipitation changes between 0 to -15°C as simulated by the Ceres-Rice model

Figure 6 shows the simulated yields and Biomass of the rice crop base on scenarios of Sra1b, Sra2 and Srb1. The Hashemi and Alikazemi rice yield decreases for 2011-2032 and

2046-2065 with temperature rise and precipitation drop. The decreasing of crop rice yields are about between 3034 to 3208 kg ha⁻¹ for Hashemi rice and 3377 to 3384 kg ha⁻¹ for Alikazemi rice.




Fig. 6. Simulation of rice yield and Biomass under three scenarios of Sra1b, Sra2 and Srb1 in the Rasht station for 2011-2032 and 2046-2065 as compared with observed data

The time of transplants and harvest will shift to winter season Because of increasing of minimum temperature. This change in recent years was obvious but this change do not recorder continuously.

Conclusion

The impacts of climate change on rice yield and Biomass will depend on the actual patterns of change in rice growing areas. The results of prediction of ECHAM5 climate model under SRA1B, SRA2 and SRB1 scenarios show that trend of temperature will be ascending in the southern coast of Caspian Sea and the minimum temperature will increase more than maximum temperature. The maximum precipitation occurs in fall season and the mean precipitation will decrease in future to the normal level. It may increase evapotranspiration level and relative humidity due to increasing of temperature. Zhiqing et al., (1994) reported that an increase in temperature alone would decrease rice yield but that enhanced photosynthesis caused by increased CO₂ can compensate for this effect. Saseendran et al., (2000) reported an increase in the trend of simulated rice yield for a 1 to 3°C drop in temperature. Aggarwall and mall (2000) show that increase of 1°C temperature without any increase in CO₂ resulted in a 3- 7% decrease in grain yield in eastern and western region India. Peng et al., (2004) concluded that grain yield declined by 10% for each 1 degrees C increase in growing-season minimum temperature in the dry season. Present study showed similar results to other researches. The sensitivity of rice yield and biomass to atmospheric temperature and precipitation changes show that for two degree temperature rise there is about 24-27% decline in simulated yield and about 26% decline in simulated biomass. These impacts can be averted through the efforts of agricultural research and policies aiming to improve rice varieties and accompanying management strategies.

References

- Aggarwall, P.K., R.K. Mall, 2002. Climate change and rice yields in diverse Agro-environments of India. II. Effect of uncertainties in scenarios and crop models on impact assessment. *Climatic Change*, 52; 331–343.
- Auffhammer, M., Ramanathan, V., J.R. Vincent, 2012. Climate change, the monsoon, and rice yield in India. *Climatic Change*, 111; 411–424.
- Bazzaz, F., W. Sombroek, 1996. Global climate change and agricultural production. Food and agriculture organization of the United Nations and John Wiley and Sons.
- Bouman, B. A. M., H. H. Van Laar, 2006. Description and evaluation of the rice growth model ORYZA2000 under nitrogen-limited conditions. *Agricultural Systems*, 87; 249–273.
- Dore, M.H.I., 2005. Climate change and changes in global precipitation patterns: What do we know?. *Environment International*, 31; 1167–1181.
- Easterling, D. R., Horton, B., D. Jones, P., Peterson, T. C., Karl, T. R., Parker, D. E., Salinger, M. J., Razuvayev, V., Plummer, N., Jamason, P., C. K. Folland, 1997. Maximum and minimum temperature trends for the globe. *Science*, 277; 364–367.
- Hewitson, B.C., R.G. Crane, 1996. Climate downscaling: techniques and application. *Climate Research*, 7; 85–95.
- Horie T., Baskar J.T., H. Nakagawa, 2000. Crop ecosystem responses to climate change: Rice. In: Reddy KR, Hodges HF, editors. *Climate change and Global crop productivity*. Wallingford: CABI Publishing; p. 81–106.
- Hunt, L.A., Pararajasingham, S., Jones, J. W., Hoogenboom, G., Imamura, D.T., R.M. Ogoshi, 1993. Gencalc: software to facilitate the use of crop models for analyzing field experiments. *Agronomy Journal*, 85; 1090–1094.
- IPCC, 2007. Climate change 2007: the physical science basis. Available from http://www.ipcc.ch/publications_and_data/publication_s_ipcc_fourth_assessment_report_wg1_report_the_physical_science_basis.htm.
- Iran Ministry of Agriculture, 1993. The feasibility study on the irrigation and drainage development project in the Haraz River Basin. Tokyo: Japan International Cooperation Agency.
- Gohari, A., Eslamian, S., Abedi-Koupaei, J., Massah Bavani, A., Wang, D., K. Madani, 2013. Climate change impacts on crop production in Iran's Zayandeh-Rud River Basin. *Science of the Total Environment*, 442; 405–419.
- Mainuddin, M., Mac Kirby, M., C.T. Hoanh, 2013. Impact of climate change on rain fed rice and options for adaptation in the lower Mekong Basin. *Nat Hazards*, 66; 905–93.
- Mathauda, S.S., Mavi, H.S., Bhangoo, B.S., B.K. Dhaliwal, 2000. Impact of projected climate change on rice production in Punjab, India. *Tropical Ecology*, 41(1); 95-98.
- Peng, S., Huang, J., Sheehy, J.E., Laza, R.C., Visperas, R.M., Zhong, X., Centeno, G.S., Khush, G.S., K.G. Cassman, 2004. Rice yields decline with higher night temperature from global warming. *Proc Natl Acad Sci USA*, 6; 9971–9975.
- Poudel, S., K. Kotani, 2013. Climatic impacts on crop yield and its variability in Nepal: do they vary across seasons and altitudes? *Climatic Change*, 116; 327–355.
- Reddy, K.R., H.F. Hodges, (Eds.), 2000. *Climate change and global crop productivity*. CABI Publishing, Wallingford, Oxon, UK.
- Ritchie, J.T., 1993. Genetic specific data for crop modeling. In: Penning de Vries FWT, Teng P, Metselaar K, editors. *Systems approaches for agricultural development*. Dordrecht: Kluwer Academic Publishers; p. 77–93.
- Saseendran, S.A., Singh, K.K., Rathore, L.S., Singh, S.V., S.K. Sinha, 2000. Effects of climate change on rice production in the tropical humid climate of Kerala, India. *Climatic Change*, 44; 495–514.
- Soora, N.K., Aggarwal, P.K., Saxena, R., Rani, S., Jain, S., N. Chauhan, 2013. An assessment of regional

- vulnerability of rice to climate change in India. *Climatic Change*, 118; 683-699.
- Timsina, J., E. Humphreys, 2006. Performance of CERES-Rice and CERES-Wheat models in rice-wheat systems: a review, *Agricultural Systems*, 90; 5-31.
- Willmott C.J., K. Matsuura, 2005. Advantages of the mean absolute error (MAE) over the root mean square error (RMSE) in assessing average model performance, *Climate Research*, 30; 79-82.
- Willmott, C.J., Ackleson, S.G., Davis, R.E., Feddema, J.J., Klink, K.M., Legates, D.R., Oconnell, J., C.M. Rowe, 1985. Statistics for the evaluation and comparison of model. *Geophys Res*, 90; 8995-9005.
- Yao, F., Xu, Y., Lin, E. Yokozawa, M., J. Zhang, 2007. Assessing the impacts of climate change on rice yields in the main rice areas of China. *Climatic Change*, 80; 395-409.
- Yao, F., Peng Cheng, Q., Zhang, J., Erd, L., B. Vijendra, 2011. Uncertainties in assessing the effect of climate change on agriculture using model simulation and uncertainty processing methods. *Atmospheric Science*, 56; 729-737.
- Yin, X., Struik, P.C., Tang, J., Qi, C., T. Liu, 2005. Model analysis of flowering phenology in recombinant inbred lines of barley. *J Exp Bot*, 56; 959-965.
- Yoshida, S., T. Hara, 1977. Effects of air temperature and light on grain filling of an Indica and Japonica rice under controlled environmental conditions. *Soil Sci. Plant Nutr*, 23; 93-107.
- Zhiqing, J., Ge, D., Chen, H., J. Fang, 1994. Effects of climate change on rice production and strategies for adaptation in southern china, In: *Implications of climate change for international agriculture: crop modeling study*, U.S. Climate Change Division Report, EPA, 230-B-94-003; p. 1-24